

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 36-1580: Monoclonal Antibody to DOG-1 / TMEM16A / ANO1 (Gastrointestinal Stromal Tumor Marker)(Clone: SPM580)

Clonality: Monoclonal Clone Name: SPM580 Application: IHC Reactivity: Human ANO1 Gene: 55107 Gene ID: **Uniprot ID:** Q5XXA6 Format: Purified

ANO1, DOG1, ORAOV2, TAOS2, TMEM16A **Alternative Name:** 

Isotype: Mouse IgG1, kappa

Immunogen Information: Recombinant human DOG-1 protein

## **Description**

Expression of DOG-1 protein is elevated in the gastrointestinal stromal tumors (GISTs), c-kit signaling-driven mesenchymal tumors of the GI tract. DOG-1 is rarely expressed in other soft tissue tumors, which, due to appearance, may be difficult to diagnose. Immunoreactivity for DOG-1 has been reported in 97.8 percent of scorable GISTs, including all c-kit negative GISTs. Overexpression of DOG-1 has been suggested to aid in the identification of GISTs, including Platelet-Derived Growth Factor Receptor Alpha mutants that fail to express c-kit antigen. The overall sensitivity of DOG1 and c-kit in GISTs is nearly identical: 94.4% vs. 94.7%.

## **Product Info**

Amount: 100 µg

Purification: Affinity Chromatography

100 µg in 500 µl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium azide is highly Content:


toxic.

Store the antibody at 4°C; stable for 6 months. For long-term storage; store at -20°C. Avoid Storage condition:

repeated freeze and thaw cycles.

## **Application Note**

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);



Formalin-fixed, paraffin-embedded human GIST stained with DOG1 Monoclonal Antibody (SPM580).