

32-20604: Recombinant Human Semaphorin 3A Fc(Discontinued)

Alternative Name : SEMA3A, SEMAD

Description

Source:CHO cells

Semaphorins are a large group of structurally-related, secreted, GPI-anchored, transmembrane, cell-signaling molecules. There are 8 major classifications of Semaphorins (the first seven ordered by number, 1-7, and the eighth designated V for virus), which are characterized by the existence of a conserved 500 amino acid SEMA domain at the amino terminus. Classes 3, 4, 6, and 7 are found in vertebrates only, whilst class 5 is found in both vertebrates and invertebrates. Each class is then divided into additional subgroups based on shared structural characteristics. Semaphorins primarily function as axon growth cone guidance factors during neuronal development. Semaphorin 3A acts as a chemo-repellent to axons, and an inhibitor of the growth of axons by signaling through receptors, Neuropilin-1 and Plexin-A. The CHO cell-derived Recombinant Human Semaphorin 3A Fc is a glycosylated, disulfide-linked homodimer of 1,976 amino acid residues, which includes the SEMA domain, immunoglobulin c2-like domain, and the C-terminal basic Arg/Lys-rich domain of the mature sequence, as well as an 8-residue N-terminal His-tag and a 230-residue C-terminal Fc region linked by two glycines. Recombinant Human Semaphorin 3A Fc has a calculated molecular weight of 226.2 kDa and therefore runs above the 200kDa marker by SDS-PAGE analysis under nonreducing conditions. When run under reducing conditions, this protein migrates as three distinct bands that, due to glycosylation, run higher than expected at apparent molecular weights of approximately 120-130 kDa, 90-100 kDa, and 35-40 kDa.

Product Info

Amount : 5 µg / 25 µg

Purification : Purity:>= 95% by SDS-PAGE gel and HPLC analyses.

Content : This recombinant protein is supplied in lyophilized form.

Amino Acid : HHHHHHHHGK NNPRLKLSY KEMLESNNVI TFNGLANSSS YHTFLLDEER SRLYVGAKDH IFSFDLVNIK
DFQKIVWPVS YTRRDECKWA GKDILKECAN FIKVLKAYNQ THLYACGTGA FHPICTYIEI GHHPEDNIFK
LENSHFENGR GKSPYDPKLL TASLLIDGEL YSGTAADFMG RDFAIFRTLH HHHPIRTEQH DSRWLNDPKF
ISAHLISESD NPEDDKVYFF FRENADIGEY SGKATHARIG QICKNDFGGH RSLVNKWTTF LKARLICSVP
GPNIDHTHFD ELQDVFLMNF KDPKNPVVYG VFTTSSNIFK GSAVCMYSMS DVRRVFLGPY
AHRDGPYQW VPYQGRVPYP RPGTCPSKTF GGFDSTKDLF DDVITFARSH PAMYNPVFPM NNRPIVIKTD
VNYQFTQIVV DRVDAEDGQY DVMFIGTDVG TVLKVVSIPK ETWYDLEEVV LEEMTVFREP TAISAMELST
KQQQLYIGST AGVAQLPLHR CDYIGKACAE CCLARDPYCA WDGSACSRYP PTAKRATRAQ DIRNGDPLTH
CSDLHHDNHH GHSPEERIIY GVENSSTFLE CSPKSQRALV YWQFQRRNEE RKEEIRVDDH IIRTDQGLLL
RSLQQKDSGN YLCHAVEHGF IQTLKVTLE VIDTEHLEEL LHKDDDGDS KTKEMSNSMT PSQKVWYRDF
MQLINHPNLN TMDEFCEQVW KRDRKQRRQR PGHTPGNSNK WKHLQENKKG RNRRTHEFER
APRSVGGPKS CDKTHTCPPC PAPELLGGPS VFLFPPKPKD TLMISRTPEV TCVVVDVSHE DPEVKFNWYV
DGVEVHNAKT KPREEQYNST YRVVSVLTVL HQDWLNGKEY KCKVSNKALP APIEKTISKA KGQPREPQVY
TLPPSRDELT KNQVSLTCLV KGFYPSDIAV EWESNGQPEN NYKTTTPVLD SDGSFFLYSK LTVDKSRWQQ
GNVFSCSVMH EALHNHYTQK SLSLSPGK

Application Note

Determined by its ability to bind recombinant rat Neuropilin-1 Fc Chimera in a functional ELISA assay.