

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

### 32-20629: Recombinant Human IFN-Beta (Discontinued)

**Reactivity:** Human, Monkey, Mouse,

Alternative Name: Fibroblast Interferon, IFNB1, Type I Interferon

# **Description**

### Source:CHO cells

Proteins of this family play an important role in inducing non-specific resistance against a broad range of viral infections. They also affect cell proliferation and modulate immune responses. Produced by peripheral blood leukocytes and lymphoblastoid cells, IFN-Alpha is an acid-stable molecule that signals through IFN-Alpha /Beta R, which is also used by IFN-Beta . Both IFNs have similar anti-viral activity and regulate expression of MHC class I antigens. IFN-Alpha contains four highly conserved cysteine residues that form two disulfide bonds, one of which is necessary for biological activity. The Recombinant Human IFN-Beta is a 20.0 kDa protein containing 166 amino acid residues. Due to glycosylation, IFN-Beta has an approximate MW of 22.3 kDa based on SDS-PAGE gel and Mass Spectrometry.

#### **Product Info**

Amount:  $5 \mu g / 20 \mu g$ 

**Purification:** Purity:>= 95% by SDS-PAGE gel and HPLC analyses. **Content:** This recombinant protein is supplied in lyophilized form.

Amino Acid: MSYNLLGFLQ RSSNFQCQKL LWQLNGRLEY CLKDRMNFDI PEEIKQLQQF QKEDAALTIY EMLQNIFAIF

RQDSSSTGWN ETIVENLLAN VYHQINHLKT VLEEKLEKED FTRGKLMSSL HLKRYYGRIL HYLKAKEYSH

CAWTIVRVEI LRNFYFINRL TGYLRN

# **Application Note**

Assay #1:Determined by its ability to induce STAT1/STAT2 activation in Human COLO 205 ISRE LUC reporter cells. Assay #2:Determined by a cytotoxicity assay using human TF-1 cells. The expected  $\hat{A} = \hat{A} = \hat{A}$