

## 32-20662: Recombinant Human sCD30 Ligand(Discontinued)

**Alternative Name :** soluble CD30 Ligand, TNFSF8, CD153, CD30L

### Description

**Source:CHO cells**

CD30 ligand (CD30L) is a type-II membrane-associated glycoprotein belonging to the TNF superfamily and is expressed primarily on certain B cells, T cells, and monocytes. CD30L binds specifically to CD30 (receptor), which is expressed on activated, but not resting, B and T cells, in lymphomas and various chronically inflamed tissues. CD30L/CD30 interactions initiate a signaling cascade that can ultimately lead to the activation of NF- $\kappa$ B. CD30L/CD30 signaling exerts pleiotropic effects on normal cells, including cell death, differentiation, and cell division. Certain diseases, including Hodgkin's lymphoma, allergic inflammation, diabetes (in NOD mice), and mycobacterial infection can also be affected by CD30L/CD30 signaling. The CD30L gene encodes for a 234 amino acid type II transmembrane protein, which contains a 37 amino acid cytoplasmic sequence, a 25 amino acid transmembrane domain and a 172 amino acid extracellular domain. Recombinant Human soluble CD30L (sCD30L) is a 188 amino acid polypeptide corresponding to the extracellular domain, and contains an 8 residue N-terminal His-Tag. The calculated molecular weight of Recombinant Human sCD30L is 21.3 kDa.

### Product Info

**Amount :** 10  $\mu$ g / 50  $\mu$ g

**Purification :** Purity: $\geq$  98% by SDS-PAGE gel and HPLC analyses.

**Content :** This recombinant protein is supplied in lyophilized form.

**Amino Acid :** HHHHHHHHPS PGGSGGQRTD SIPNSPDNVP LKGGNCSEDL LCILKRPFK KSWAYLQVAK  
HLNKTCLSWN KDGILHGVRV QDGNLVIQFP GLYFIICQLQ FLVQCPNNSV DLKLELLINK HIKKQALVTV  
CESGMQTKHV YQNLSQFLLD YLQVNTTISV NVDTFQYIDT STFPLENVLS IFLYSNSD

### Application Note

Determined by its ability to stimulate human IL-8 production by human PBMC using a concentration range of 10.0-25.0 ng/ml.