w abeomics

37-1041: Human ERK2 / MAPK1 / MAPK2 Recombinant Protein (GST Tag)(Discontinued)

Reactivity : Human

Alternative Name : ERK Protein, ERK-2 Protein, ERK2 Protein, ERT1 Protein, MAPK2 Protein, p38 Protein, p40 Protein, p41 Protein, p41mapk Protein, p42-MAPK Protein, P42MAPK Protein, PRKM1 Protein, PRKM2 Protein, PA

Description

Source : Baculovirus-Insect Cells

MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. ERK is a versatile protein kinase that regulates many cellular functions. Growing evidence suggests that extracellular signal-regulated protein kinase 1/2 (ERK1/2) plays a crucial role in promoting cell death in a variety of neuronal systems, including neurodegenerative diseases. It is believed that the magnitude and the duration of ERK1/2 activity determine its cellular function. Activation of ERK1/2 are implicated in the pathophysiology of spinal cord injury (SCI). ERK2 signaling is a novel target associated with the deleterious consequences of spinal injury. ERK-2, also known as Mitogenactivated protein kinase 1 (MAPK1), is a member of the protein kinase superfamily and MAP kinase subfamily. MKP-3 is a dual specificity phosphatase exclusively specific to MAPK1 for its substrate recognition and dephosphorylating activity. The activation of MAPK1 requires its phosphorylation by upstream kinases. Upon activation, MAPK1 translocates to the nucleus of the stimulated cells, where it phosphorylates nuclear targets. MAPK1 is involved in both the initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by phosphorylating a number of transcription factors such as ELK1. MAPK1 acts as a transcriptional repressor which represses the expression of interferon gamma-induced genes. Transcriptional activity is independent of kinase activity. The nuclear-cytoplasmic distribution of ERK2 is regulated in response to various stimuli and changes in cell context. Furthermore, the nuclear flux of ERK2 occurs by several energy- and carrier-dependent and -independent mechanisms. ERK2 has been shown to translocate into and out of the nucleus by facilitated diffusion through the nuclear pore, interacting directly with proteins within the nuclear pore complex, as well as by karyopherin-mediated transport. ERK2 interacts with the PDE4 catalytic unit by binding to a KIM (kinase interaction motif) docking site located on an exposed beta-hairpin loop and an FQF (Phe-GIn-Phe) specificity site located on an exposed alphahelix. These flank a site that allows phosphorylation by ERK, the functional outcome of which is orchestrated by the Nterminal UCR1/2 (upstream conserved region 1 and 2) modules. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

Product Info

Amount :	Human ERK2 / MAPK1 / MAPK2 Recombinant Protein (GST Tag)(Discontinued) / 50 µg		
Purification :	> 98 % as determined by SDS-PAGE		
Content :	Formulation Lyophilized from sterile 50mM Tris, 100mM NaCl, 0.5mM PMSF, 10% Glycerol, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.		
Storage condition :	Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.		
Amino Acid :	Met1-Ser360		

Application Note

Endotoxin :< 1.0 EU per μg of the protein as determined by the LAL method



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com

KDa	м	
116	-	
66.2		-
45.0	-	
35.0	-	
25.0		
18,4	_	
14.4	_	

Fig 1: Human ERK2 / MAPK1 / MAPK2 Recombinant Protein (GST Tag)