

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

37-1044: Human CD153 / CD30L / TNFSF8 Recombinant Protein (Fc Tag)(Discontinued)

Reactivity: Human

Alternative Name: CD153 Protein, CD30L Protein, CD30LG Protein,

Description

Source: HEK293 Cells

CD3 ligand (CD3L), also known as CD153 and TNFSF8, is a membrane-associated glycoprotein belonging to the TNF superfamily and TNFR superfamily, and is a specific ligand for CD3/TNFRSF8 originally described as a cell surface antigen and a marker for Hodgkin lymphoma and related hematologic malignancies. CD3L is a type-II membrane glycoprotein expressed on activated T cells, stimulated monocyte-macrophages, granulocytes, eosinophils, and some Burkitt-like lymphoma cell lines. CD3L is capable of transducing signals through CD3 on different CD3+ lymphoma cell lines, and mediates pleiotropic biologic effects including cell proliferation, activation, differentiation, as well as cell death by apoptosis. CD3-CD3 ligand interaction has been suggested to have a pathophysiologic role in malignant lymphomas, particularly Hodgkin disease, large cell anaplastic lymphomas and Burkitt lymphomas, and is also involved in activation and functioning of the T cell-dependent immune response. Thus, CD153 and its receptor CD3 are regarded as therapeutic targets in hematologic malignancies, autoimmune and inflammatory diseases.

Product Info

Amount: Human CD153 / CD30L / TNFSF8 Recombinant Protein (Fc Tag)(Discontinued) / 50 µg

Purification: > 95 % as determined by SDS-PAGE

Formulation Lyophilized from sterile PBS, pH 7.4

Content: Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

Storage condition:

Storage condition:

Storage condition:

Storage condition:

for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Gln63-Asp234

Application Note

Measured by its binding ability in a functional ELISA. Immobilized recombinant human CD30L at 20 μ g/ml (100 μ L/well) can bind biotinylated human CD30 with a linear range of 0.31-20 ng/ml .

Endotoxin :< 1.0 EU per μg of the protein as determined by the LAL method.

Other pack size also available.




Fig 1: Human CD153 / CD30L / TNFSF8 Recombinant Protein (Fc Tag)