

37-1072: Human CD146 / MCAM Recombinant Protein (Fc Tag)(Discontinued)

 Reactivity :
 Human

 Alternative Name :
 CD146 Protein, MUC18 Protein,

Description

Source : HEK293 Cells

The CD146 antigen, also known as melanoma cell adhesion molecule (MCAM) and MUC18, is an integral membrane glycoprotein belonging to the immunoglobulin superfamily. CD146 contains the characteristic immunoglobulin-like domains (V-V-C2-C2-C2), a transmembrane region and a short cytoplasmic tail. The CD146 expression is detected in endothelial cells in vascular tissue throughout the body, and plays a role in cell adhesion, as well as in cohesion of the endothelial monolayer at intercellular junctions in vascular tissue. As a Ca2+-independent cell adhesion molecule involved in heterophilic cell to cell interactions and a surface receptor, CD146 triggers tyrosine phosphorylation of FYN and PTK2 and subsequently induced signal transduction, proteolysis, or immune recognition. This protein is also expressed predominantly on metastatic lesions and advanced primary tumours, and thus has been suggested to play an important role in tumour progression and the development of metastasis in certain human carcinomas.

Product Info

Amount :	Human CD146 / MCAM Recombinant Protein (Fc Tag)(Discontinued) / 50 µg
Purification :	> 95 % as determined by SDS-PAGE
Content :	Formulation Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.
Storage condition :	Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Amino Acid :	Met1-Gly559
Application Note	

Endotoxin :< 1.0 EU per μ g of the protein as determined by the LAL method

KDa	M		
116	-	-	-
66.2	-		
45.0	-		
35.0	-		
25.0	-		
18.4	-		
14.4	-		

Fig 1: Human CD146 / MCAM Recombinant Protein (Fc Tag)

For Research Use Only. Not for use in diagnostic/therapeutics procedures.