

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

37-1137: Human HIF-1 alpha / HIF1A Recombinant Protein (His Tag)(Discontinued)

Reactivity: Human

Alternative Name: bHLHe78 Protein, HIF-1A Protein, HIF-1alpha Protein, HIF1 Protein, HIF1-ALPHA Protein, MOP1 Protein, HIF-1alpha Protein, HIF1-ALPHA Protein, MOP1 Protein, HIF1-ALPHA Protein, MOP1 Protein, HIF1-ALPHA Protein, MOP1 Protein, HIF1-ALPHA Protein, HIF1-ALPHA Protein, MOP1 Protein, HIF1-ALPHA Protein, HIF1-ALPHA Protein, MOP1 Protein, HIF1-ALPHA Protein, HIF1-

PASD8 Protein.

Description

Source: E. coli

HIF-1 alpha, also known as HIF1A, contains 1 basic helix-loop-helix (bHLH) domain, 1 PAC (PAS-associated C-terminal) domain and 2 PAS (PER-ARNT-SIM) domains. It is one of the two subunits of Hypoxia-inducible factor-1 (HIF1). HIF1 is a transcription factor found in mammalian cells cultured under reduced oxygen tension that plays an essential role in cellular and systemic homeostatic responses to hypoxia. HIF1 is a heterodimer composed of an alpha subunit and a beta subunit. The beta subunit has been identified as the aryl hydrocarbon receptor nuclear translocator (ARNT). HIF-1 alpha is expressed in most tissues with highest levels in kidney and heart. It is overexpressed in the majority of common human cancers and their metastases, due to the presence of intratumoral hypoxia and as a result of mutations in genes encoding oncoproteins and tumor suppressors. HIF-1 alpha functions as a master transcriptional regulator of the adaptive response to hypoxia. Under hypoxic conditions, it activates the transcription of over 4 genes, including erythropoietin, glucose transporters, glycolytic enzymes, vascular endothelial growth factor, HILPDA, and other genes whose protein products increase oxygen delivery or facilitate metabolic adaptation to hypoxia. HIF1A plays an essential role in embryonic vascularization, tumor angiogenesis and pathophysiology of ischemic disease. HIF-1 alpha binds to core DNA sequence 5'-[AG]CGTG-3' within the hypoxia response element (HRE) of target gene promoters. Activation requires recruitment of transcriptional coactivators such as CREBPB and EP3.

Product Info

Amount: 1 alpha / HIF1A Recombinant Protein (His Tag)(Discontinued) / 100 µg

Purification: > 95 % as determined by SDS-PAGE

Formulation Lyophilized from sterile PBS, pH 7.4

Content: Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

Storage condition:

Storage condition:

Storage condition:

Storage condition:

for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Arg575-Asn826




Fig 1: Human HIF-1 alpha / HIF1A Recombinant Protein (His Tag)