

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

37-1189: Mouse CD160 Recombinant Protein (Fc Tag)(Discontinued)

Reactivity: Mouse

Alternative Name: AU045688 Protein, Mouse; By55 Protein, Mouse

Description

Source: HEK293 Cells

CD16 antigen, also known as Natural killer cell receptor BY55 and CD16, is a cell membrane protein which contains one Iglike V-type (immunoglobulin-like) domain. CD16 is a GPI-anchored lymphocyte surface receptor in which expression is mostly restricted to the highly cytotoxic CD56(dim)CD16(+) peripheral blood NK subset. CD16 is a receptor showing broad specificity for both classical and non-classical MHC class I molecules. CD16 is expressed in spleen, peripheral blood, and small intestine. Expression of CD16 is restricted to functional NK and T cytotoxic lymphocytes. CD16 acts as a co-activator receptor for CD3-induced proliferation of CD4+ CD16+ T cells isolated from inflammatory skin lesions. Unique CD4+ CD16+ lymphocyte subset may play a role in the pathogenesis of skin inflammation. Activated NK lymphocytes release a soluble form of CD16 that functionally impairs the MHC-I-specific cytotoxic CD8(+) T lymphocyte responsiveness. Cancer Immunotherapy Co-inhibitory Immune Checkpoint Targets Immune Checkpoint Immune Checkpoint Detection: Antibodies Immune Checkpoint Detection: FCM Antibodies Immune Checkpoint Detection: IHC Antibodies Immune Checkpoint Targets Immunotherapy Targeted Therapy

Product Info

Amount: Mouse CD160 Recombinant Protein (Fc Tag)(Discontinued) / 200 µg

Purification: > 95 % as determined by SDS-PAGE.

Formulation Lyophilized from sterile PBS, pH 7.4.

Content: Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

Storage condition : Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be

aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Met1-Leu159

Application Note

Endotoxin :< 1.0 EU per ̸µg protein as determined by the LAL method.




Fig 1: Mouse CD160 Recombinant Protein (Fc Tag)