

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

37-1218: Human ICOS / AILIM / CD278 Recombinant Protein (Fc Tag)(Discontinued)

Reactivity: Human

Alternative Name: AILIM Protein, CD278 Protein, CVID1 Protein,

Description

Source: HEK293 Cells

Inducible costimulator (ICOS), also called AILIM (activiation-inducible lymphocyte immunomediatory molecule) is a cell-surface receptor, and belongs to the CD28 family of immune costimulatory receptors consisting of CD28, CTLA-4 and PD-1. The interaction of B7-H2/ICOS plays a critical role in Th cell differentiation, T-B cell interactions which is essential for germinal center formation, and humoral immune responses, and as well as the production of cytokine IL-4. In addition, ICOS is more potent in the induction of IL-1 production, a cytokine important for suppressive function of T regulatory cells. The B7-1/B7-2-CD28/CTLA-4 and ICOS-B7RP-1 pathway provides key second signals that can regulate the activation, inhibition and fine-tuning of T-lymphocyte responses. ICOS stimulates both Th1 and Th2 cytokine production but may have a preferential role in Th2 cell development. Moreover, The B7-1/B7-2-CD28/CTLA-4 and ICOS-B7RP-1 pathway has been suggested of being involved in the development of airway inflammation and airway hyperresponsiveness. Cancer Immunotherapy Co-stimulatory Immune Checkpoint Targets Immune Checkpoint Immune Checkpoint Proteins Immune Checkpoint Targets Immunotherapy Targeted Therapy

Product Info

Amount: Human ICOS / AILIM / CD278 Recombinant Protein (Fc Tag)(Discontinued) / 100 µg

Purification: > 95 % as determined by SDS-PAGE.

Formulation Lyophilized from sterile PBS, pH 7.4.

Content: Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

Storage condition:

Storage condition:

Storage condition:

Storage condition:

for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Met1-Phe141

Application Note

Endotoxin :< 1.0 EU per µg protein as determined by the LAL method.




Fig 1: Human ICOS / AILIM / CD278 Recombinant Protein (Fc Tag)