

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

# 37-1224: Human TIMP2 / TIMP-2 Recombinant Protein (Fc Tag)(Discontinued)

**Reactivity:** Human

Alternative Name: CSC-21K Protein, DDC8 Protein,

## **Description**

#### Source: HEK293 Cells

Tissue inhibitors of metalloproteinases (TIMP) family are natural inhibitors of the matrix metalloproteinases (MMPs), the zinc enzymes involved in extracellular matrix maintenance and remodeling. The TIMP family encompasses four members (TIMP1-4), and they inhibit most MMPs by forming non-covalent binary complex. TIMP2 is a 22 kDa non N-glycosylated protein expressed by a variety of cell types, and plays a unique role among TIMP family members owing to its functions to regulate cellular responses to growth factors. Findings establish an unexpected, MMP-independent mechanism for TIMP2 inhibition of endothelial cell proliferation in vitro and reveal an important component of the antiangiogenic effect of TIMP2 in vivo. TIMP-2 thus is critical to the maintenance of tissue homeostasis and is involved in the regulation of tumor microenvironment.

### **Product Info**

**Amount :** 2 Recombinant Protein (Fc Tag)(Discontinued) / 20 μg

**Purification:** > 95 % as determined by SDS-PAGE

Formulation Lyophilized from sterile 100mM Glycine, 10mM NaCl, 50mM Tris, pH 7.5

**Content:** Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

**Storage condition :** Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be

aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Cys27-Pro220

### **Application Note**

Endotoxin :< 1.0 EU per ̸µg of the protein as determined by the LAL method




Fig 1: Human TIMP2 / TIMP-2 Recombinant Protein (Fc Tag)