

## 37-1288: Human XPNPEP2 / X-Pro aminopeptidase 2 Recombinant Protein (His Tag)(Discontinued)

**Reactivity :** Human

**Alternative Name :** AEACEI Protein, APP2 Protein,

### Description

#### Source : HEK293 Cells

Aminopeptidase P (APP) is a hydrolase specific for N-terminal imido bonds, which are common to several collagen degradation products, neuropeptides, vasoactive peptides, and cytokines. A membrane-bound and soluble form of this enzyme (XPNPEP2) have been identified as products of two separate genes. XPNPEP2, the X-linked gene that encodes membranous aminopeptidase P (APP), has been reported to associate with APP activity. The membrane aminopeptidase P (XPNPEP2) is largely limited in distribution to endothelia and brush border epithelia. APP and XPNPEP2 contain homologous blocks of sequence common to members of the "pita bread-fold" protein family, of which Escherichia coli methionine aminopeptidase is the prototype. The C-2399A variant in XPNPEP2 is associated with reduced APP activity and a higher incidence of AE-ACEi. XPNPEP2 mRNA was detected in fibroblasts that carry the translocation, suggesting that this gene at least partially escapes X inactivation. XPNPEP2 is a candidate gene for premature ovarian failure (POF).

### Product Info

**Amount :** Pro aminopeptidase 2 Recombinant Protein (His Tag)(Discontinued) / 20 µg

**Purification :** > 97 % as determined by SDS-PAGE

Formulation Lyophilized from sterile PBS, pH 7.4

**Content :** Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.

**Storage condition :** Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

**Amino Acid :** Met1-Ala650

### Application Note

Measured by its ability to cleave the fluorogenic peptide substrate, H-Lys(2-Aminobenzoyl)Pro-Pro-pNitroanilide(K(Abz)PP-pNA). The specific activity is > 300 pmoles/min/Åµg.

Endotoxin :< 1.0 EU per Åµg of the protein as determined by the LAL method




Fig 1: Human XPNPEP2 / X-Pro aminopeptidase 2 Recombinant Protein (His Tag)