

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

# 37-1301: Human OX-40L / TNFSF4 / CD252 Recombinant Protein (Fc Tag)(Discontinued)

**Reactivity:** Human

Alternative Name: CD134L Protein, CD252 Protein, GP34 Protein, OX-40L Protein, OX40L Protein, TXGP1 Protein,

# **Description**

#### Source: HEK293 Cells

OX-4L, also known as TNFSF4 and CD252, is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. OX-4L is an important costimulatory molecule that plays a crucial role in the regulation of T-cell-mediated immunity. The interaction of TNFSF4-TNFSF4 is involved in the pathogenesis of multiple autoimmune and inflammatory diseases such as systemic lupus erythematosus (SLE), carotid artery disease and cancer. OX-4L is a ligand for receptor TNFRSF4/OX4. It is found to play a role in T cell antigen-presenting cell (APC) interactions. In surface Ig- and CD4-stimulated B cells, this cytokine along with CD7 has been shown to provide CD28-independent costimulatory signals to T cells. This protein and its receptor are reported to directly mediate adhesion of activated T cells to vascular endothelial cells. Cancer Immunotherapy Co-stimulatory Immune Checkpoint Targets Immuno Checkpoint Targets Immunotherapy Targeted Therapy

### **Product Info**

Amount: 40L / TNFSF4 / CD252 Recombinant Protein (Fc Tag)(Discontinued) / 100 μg

**Purification:** > 90 % as determined by SDS-PAGE.

Formulation Lyophilized from sterile PBS, pH 7.4.

**Content:** Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

Storage condition:

Storage condition:

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be

aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Gln51-Leu183

## **Application Note**

Measured by its binding ability in a functional ELISA.Immobilized recombinant Human TNFRSF4-His at 10  $\tilde{A} \Box \hat{A} \mu g/ml$  (100  $\tilde{A} \Box \hat{A} \mu L/well$ ) can bind human S4-mFc3-TNFSF4 with a linear range of 1.28-20  $\tilde{A} \Box \hat{A} \mu g/ml$ .Endotoxin :< 1.0 EU per  $\tilde{A} \Box \hat{A} \mu g/ml$  protein as determined by the LAL method.




Fig 1: Human OX-40L / TNFSF4 / CD252 Recombinant Protein (Fc Tag)



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig 2: Human OX-40L / TNFSF4 / CD252 Recombinant Protein (Fc Tag) measured by its binding ability in a functional ELISA.