

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

37-1340: Mouse CXCL2 / MIP-2 Recombinant Protein (His Tag)(Discontinued)

Reactivity: Mouse

CINC-2a Protein, Mouse; Gro2 Protein, Mouse; GROb Protein, Mouse; Mgsa-b Protein, Mouse; MIP-2

Alternative Name: Protein, Mouse; MIP-2a Protein, Mouse; Mip2 Protein, Mouse; Scyb Protein, Mouse; Scyb2 Protein,

Mouse

Description

Source: Yeast

Chemokine (C-X-C motif) ligand 2 (CXCL2), also called macrophage inflammatory protein 2 (MIP-2), Growth-regulated protein beta (Gro-beta) and Gro oncogene-2 (Gro-2), is a small cytokine belonging to the CXC chemokine family. CXCL2/MIP-2 is selectively up-regulated in tolerance-conferring APCs and serves to recruit NKT cells to the splenic marginal zone, where they form clusters with APCs and T cells. In the absence of the high-affinity receptor for CXCL2/MIP-2 or in the presence of a blocking Ab to CXCL2/MIP-2, peripheral tolerance is prevented, and Ag-specific T regulatory cells are not generated. CXCL2/MIP-2 is selectively up-regulated in tolerance-conferring APCs and serves to recruit NKT cells to the splenic marginal zone, where they form clusters with APCs and T cells. In the absence of the high-affinity receptor for MIP-2 (as in CXCR2deficient mice) or in the presence of a blocking Ab to MIP-2, peripheral tolerance is prevented, and Ag-specific T regulatory cells are not generated. Understanding the regulation of lymphocyte traffic during tolerance induction may lead to novel therapies for autoimmunity, graft acceptance, and tumor rejection. Several studies have implicated the CXCL2 chemokine as a mediator in the development of sepsis. CXCL2/MIP-2 also plays a major role in mediating the neutrophilic inflammatory response of the rodent lung to particles such as quartz, crocidolite asbestos, as well as high doses of other relative innocuous dusts such as titanium dioxide.

Product Info

Amount: 2 Recombinant Protein (His Tag)(Discontinued) / 50 μg

Purification: > 95 % as determined by SDS-PAGE.

Formulation Lyophilized from sterile 30% CAN.

Content: Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted Storage condition:

for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Ala28-Asn100

Application Note

Other pack size also available




Fig 1: Mouse CXCL2 / MIP-2 Recombinant Protein (His Tag)