

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

37-1369: Mouse EphA1 / EPH receptor A1 Recombinant Protein (His Tag)(Discontinued)

Reactivity: Mouse

Alternative Name: 5730453L17Rik Protein, Mouse; AL033318 Protein, Mouse; Eph Protein, Mouse; Esk Protein, Mouse

Description

Source: HEK293 Cells

EPHA1 or EPH receptor A1 belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. An important role of Eph receptors and their ligands ephrins is to mediate cell-contact-dependent repulsion. Eph receptors and ephrins also act at boundaries to channel neuronal growth cones along specific pathways, restrict the migration of neural crest cells, and via bidirectional signaling prevent intermingling between hindbrain segments. Eph receptors and ephrins can also trigger an adhesive response of endothelial cells and are required for the remodeling of blood vessels. Eph receptors and ephrins have emerged as key regulators of the repulsion and adhesion of cells that underlie the establishment, maintainence, and remodeling of patterns of cellular organization. The ephrins and Eph receptors are implicated as positional labels that may guide the development of neural topographic maps.

Product Info

Amount: Mouse EphA1 / EPH receptor A1 Recombinant Protein (His Tag)(Discontinued) / 200 μg

Purification: > 97 % as determined by SDS-PAGE

Formulation Lyophilized from sterile PBS, pH 7.4

Content: Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

Storage condition:

Storage condition:

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be

aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Met1-Glu548

Application Note

Measured by its binding ability in a functional ELISA. Immobilized mouse EPHA1-His at 10 $\tilde{A} \Box \hat{A} \mu g/ml$ (100 $\tilde{A} \Box \hat{A} \mu L/well$) can bind mouse EFNA1-Fc , The EC50 of mouse EFNA1-Fc is 21.3-49.8 ng/ml.

Endotoxin :< 1.0 EU per $\tilde{A} \square \hat{A} \mu g$ of the protein as determined by the LAL method




Fig 1: Mouse EphA1 / EPH receptor A1 Recombinant Protein (His Tag)