

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

37-1374: Mouse EphA3 Recombinant Protein (Fc Tag)(Discontinued)

Reactivity: Mouse

Alternative Name: AW492086 Protein, Mouse; Cek4 Protein, Mouse; End3 Protein, Mouse; ETK1 Protein, Mouse; Hek

Protein, Mouse; Hek4 Protein, Mouse; Mek4 Protein, Mouse; Tyro4 Protein, Mouse

Description

Source: HEK293 Cells

EPHA3 gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH and EPH-related receptors have been implicated in mediating developmental events, particularly in the nervous system. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. EPHA3 gene encodes a protein that binds ephrin-A ligands. EPHA3 is involved in the retinotectal mapping of neurons. It may also control the segregation but not the guidance of motor and sensory axons during neuromuscular circuit development. Cancer Immunotherapy Immune Checkpoint Immunotherapy Targeted Therapy

Product Info

Amount: Mouse EphA3 Recombinant Protein (Fc Tag)(Discontinued) / 500 µg

Purification: > 90 % as determined by SDS-PAGE

Formulation Lyophilized from sterile PBS, pH 7.4.

Content: Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

Storage condition:

Storage condition:

Storage condition:

Storage condition:

Storage condition:

for optimal storage. Avoid repeated freeze-thaw cycles.

Amino Acid: Met1-His541

Application Note

Measured by its binding ability in a functional ELISA. Immobilized mouse EFNA5-His at 10 $\text{Å}\mu\text{g/ml}$ (100 $\text{Å}\mu\text{L/well}$) can bind mouse EPHA3-Fc, The EC50 of rat mouse EPHA3-Fc is 12.3-28.9 ng/ml. Endotoxin :< 1.0 EU per $\text{Å}\mu\text{g}$ of the protein as determined by the LAL method




Fig 1: Mouse EphA3 Recombinant Protein (Fc Tag)