

36-2057: Anti-LH, alpha (Luteinizing Hormone, alpha) Monoclonal Antibody (Clone: LHa/756)

Clonality :	Monoclonal
Clone Name :	LHa/756
Application :	IHC
Reactivity :	Human
Gene :	LHA
Gene ID :	1081
Uniprot ID :	P01215
Alternative Name :	Chorionic Gonadotrophin Alpha (CGA); Follicle Stimulating Hormone Alpha (FSHA); Follicle Stimulating Hormone Alpha; GPH Alpha; GPHA1; LH-alpha; Luteinizing Hormone Alpha (LHA); Lutropin Alpha; Thyroid Stimulating Hormone Alpha (TSHA); Thyrotropin Alpha;
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Recombinant full-length hLH alpha protein

Description

This MAb reacts with a protein of ~13kDa, identified as alpha sub-unit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunits vary. LH has a beta subunit of 121 amino acids (LHB) that confers its specific biologic action and is responsible for interaction with the LH receptor. This beta subunit contains the same amino acids in sequence as the beta subunit of hCG and both stimulate the same receptor; however, the hCG beta subunit contains an additional 24 amino acids and the hormones differ in the composition of their sugar moieties. LH is synthesized and secreted by gonadotrophs in the anterior lobe of the pituitary gland. In concert with the other pituitary gonadotropin follicle-stimulating hormone (FSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary tumors and the study of pituitary disease.

Product Info

Amount :	20 µg / 100 µg
Content :	200µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(No special pre-treatment is required for staining of formalin-fixed tissues. Note that HIER is NOT recommended for this antibody.)




Fig.1: Formalin-fixed, paraffin-embedded human Pituitary stained with LH alpha Mouse Monoclonal Antibody (LHa/756).