

## 36-2099: Anti-Adipophilin / Perilipin-2 (Marker of Lipid Accumulation) Monoclonal Antibody (Clone: ADFP/1494)

|                                |  |
|--------------------------------|--|
| <b>Clonality :</b>             | Monoclonal   |
| <b>Clone Name :</b>            | ADFP/1494  |
| <b>Application :</b>           | ELISA, IHC   |
| <b>Reactivity :</b>            | Human  |
| <b>Gene :</b>                  | PLIN2  |
| <b>Gene ID :</b>               | 123  |
| <b>Uniprot ID :</b>            | Q99541   |
| <b>Alternative Name :</b>      | Adipophilin; ADFP; Adipose differentiation-related protein (ADRP); Perilipin-2 (PLIN2)                       |
| <b>Isotype :</b>               | Mouse IgG2b, kappa   |
| <b>Immunogen Information :</b> | Recombinant fragment (around aa 249-376) of human Adipophilin (ADFP) protein (exact sequence is proprietary) |

### Description

Recognizes a protein of 48kDa, which is identified as Adipophilin. It belongs to the perilipin family, members of which coat intracellular lipid storage droplets. This protein is associated with the lipid globule surface membrane material, and maybe involved in development and maintenance of adipose tissue. However, it is not restricted to adipocytes as previously thought, but is found in a wide range of cultured cell lines, including fibroblasts, endothelial and epithelial cells, and tissues, such as lactating mammary gland, adrenal cortex, Sertoli and Leydig cells, and hepatocytes in alcoholic liver cirrhosis, suggesting that it may serve as a marker of lipid accumulation in diverse cell types and diseases.

### Product Info

|                            |  |
|----------------------------|--|
| <b>Amount :</b>            | 20 µg / 100 µg   |
| <b>Content :</b>           | 200µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| <b>Storage condition :</b> | Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months.   |

### Application Note

ELISA (For coating, order Ab without BSA); Immunohistochemistry (Formalin-fixed) (2-4ug/ml for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig.1: Formalin-fixed, paraffin-embedded human Adrenal stained with Adipophilin Mouse Monoclonal Antibody (ADFP/1494).




Fig. 2: SDS-PAGE Analysis Purified Adipophilin Mouse Monoclonal Antibody (ADFP/1494). Confirmation of Integrity and Purity of Antibody.