

## 36-2124: Anti-Granulocyte-Colony Stimulating Factor (G-CSF) Monoclonal Antibody(Clone: CSF3/3166R)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	CSF3/3166R
<b>Application :</b>	IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	CSF3
<b>Gene ID :</b>	1440
<b>Uniprot ID :</b>	P09919
<b>Alternative Name :</b>	Pluripoietin; Filgrastim; Lenograstim; CSF3OS; G-CSF; chromosome 17 open reading frame 33 (C17orf33); colony stimulating factor 3 (granulocyte)
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Recombinant full-length human CSF3 protein

### Description

This MAb recognizes granulocyte-colony stimulating factor (G-CSF) in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell types. Markers of myeloid cells are useful in the identification of different levels of cellular differentiation. It reacts with early precursor and mature forms of myeloid cells. It is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes. G-CSF is a pleiotropic cytokine that influences differentiation, proliferation and activation of the neutrophilic granulocyte lineage. The human G-CSF cDNA encodes a 207 amino acid precursor containing a 29 amino acid signal peptide that is proteolytically cleaved to form a 178 amino acid residue mature protein. Two G-CSF s, which are identical except for a three amino acid deletion in the amino-terminus of one form of the protein have been isolated from human cells. Murine and human G-CSF s share 73% sequence identity at the amino acid level.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) ,(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil stained with G-CSF Recombinant Rabbit Monoclonal Antibody (CSF3/3166R).




Fig. 2: SDS-PAGE Analysis Purified G-CSF Recombinant Rabbit Monoclonal Antibody (CSF3/3166R). Confirmation of Purity and Integrity of Antibody.