

### 36-2136: Anti-Catenin, beta (p120) Monoclonal Antibody(Clone: 6F9)

|                                |  |
|--------------------------------|--|
| <b>Clonality :</b>             | Monoclonal   |
| <b>Clone Name :</b>            | 6F9  |
| <b>Application :</b>           | FACS,IF,WB,IHC   |
| <b>Reactivity :</b>            | Human  |
| <b>Gene :</b>                  | CTNNB1   |
| <b>Gene ID :</b>               | 1499   |
| <b>Uniprot ID :</b>            | P35222   |
| <b>Alternative Name :</b>      | Cadherin associated protein, beta 1 88kDa, Catenin beta-1, CATNB, CHBCAT, CTNNB1 |
| <b>Isotype :</b>               | Mouse IgG1, kappa  |
| <b>Immunogen Information :</b> | Recombinant chicken $\beta$ -catenin   |

#### Description

This MAb recognizes a protein of 92kDa, which is identified as beta-catenin. It shows no cross-reaction with gamma-catenin (also known as plakoglobin). The catenins, alpha, beta and gamma bind to the highly conserved, intracellular cytoplasmic tail of E-cadherin. Together, the catenin/cadherin complexes play an important role mediating cellular adhesion. Alpha-catenin was initially described as an E-cadherin-associated protein, and has been shown to associate with other members of the cadherin family, such as N-cadherin and P-cadherin. Beta-catenin associates with the cytoplasmic portion of E-cadherin, which is necessary for the function of E-cadherin as an adhesion molecule. Beta-catenin has also been found in complexes with the tumor suppressor protein APC.

#### Product Info

|                            |  |
|----------------------------|--|
| <b>Amount :</b>            | 20 $\mu$ g / 100 $\mu$ g   |
| <b>Content :</b>           | 200 $\mu$ g/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| <b>Storage condition :</b> | Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.                                    |

#### Application Note

Flow Cytometry (1-2 $\mu$ g/million cells); Immunofluorescence (1-2 $\mu$ g/ml); Western Blot (1-2 $\mu$ g/ml); Immunohistochemistry (Formalin-fixed) (1-2 $\mu$ g/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 9.0, for 45 min at 95 $\pm$ degC followed by cooling at RT for 20 minutes)




Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil stained with Beta-Catenin (p120) Monoclonal Antibody (6F9).




Fig. 2: Confocal immunofluorescence image of HeLa cells using Beta-Catenin (p120) Monoclonal Antibody (6F9). Green (CF488) and Reddot is used to label the nuclei.




Fig. 3: SDS-PAGE Analysis of Purified Beta-Catenin (p120) Monoclonal Antibody (6F9). Confirmation of Purity and Integrity of Antibody