

36-2171: Anti-Desmocollin-2/3 Monoclonal Antibody(Clone: 7G6)

Clonality :	Monoclonal
Clone Name :	7G6
Application :	FACS,IF,WB,IHC
Reactivity :	Human, Mouse, Rat
Gene :	DSC2
Gene ID :	1824
Uniprot ID :	Q02487
Alternative Name :	ARVD11; Cadherin family member 2; CDHF2; Desmocollin-2; Desmocollin-3; Desmosomal glycoprotein II and III; Desmosomal glycoprotein II; Desmosomal glycoprotein III; DG2; DGI/III; DSC2; DSC3
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Human desmocollin-2 extracellular domain

Description

Desmosomes are intercellular adhering junctions that represent cell surface attachment sites for intermediate filament. The desmosome is subdivided into two regions. The plaque region lies adjacent to the plasma, and is believed to contain molecules that attach the intermediate filament cytoskeleton to the desmosome. The core region is composed of transmembrane glycoproteins that are thought to mediate cell-cell adhesion. Desmogleins and desmocollins are the main desmosomal transmembrane proteins. These desmosomal glycoproteins belong to the members of the cadherin family of adhesion molecules. Three different isoforms of both desmogleins and desmocollins have been identified, named as desmoglein 1-3 and desmocollins. Desmosomal cadherins showed differentiation-specific expression in the human epidermis, although the functional significance of this differential expression is not fully understood. Desmocollin-1 can be found in the upper layers. The expression of desmocollin-2 varies in the basal and suprabasal layers. And desmocollin-3 is expressed more evenly throughout the suprabasal layers.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Western Blot (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes)




Fig. 1: Formalin-fixed, paraffin-embedded human Skin stained with Desmocollin-2/3 Mouse Monoclonal Antibody (7G6).




Fig. 2: Formalin-fixed, paraffin-embedded human Skin stained with Desmocollin-2/3 Mouse Monoclonal Antibody (7G6).




Fig. 3: SDS-PAGE Analysis Purified Desmocollin-2/3 Mouse Monoclonal Antibody (7G6). Confirmation of Purity and Integrity of Antibody.