w abeomics

36-2174: Anti-Desmoglein-1 (DSG1) Monoclonal Antibody(Clone: 27B2)

Clonality :	Monoclonal
Clone Name :	27B2
Application :	FACS,IF,WB
Reactivity :	Human
Gene :	DSG1
Gene ID :	1828
Uniprot ID :	Q02413
Alternative Name :	Cadherin family member 4; CDHF4; Desmoglein-1; Desmosomal glycoprotein 1; DG1; DSG1; EPKHE; EPKHIA; Pemphigus foliaceus antigen; PPKS1; SPPK1
Isotype :	Mouse IgG1, kappa
Immunogen Information	: Human desmoglein-1 intracellular domain

Description

Recognizes a protein of ~150kDa, identified as Desmoglein-1 (DSG1). Desmoglein-1 is a member of the desmosomal cadherin family.Desmosomes are intercellular adhering junctions that represent cell surface attachment sites for intermediate filament. Desmocollins and desmogleins are the main desmosomal transmembrane proteins. Desmogleins consist of Dsg1, Dsg2, Dsg3, and Dsg4 isoforms.Within the desmosome, the extracellular domain of desmoglein is essential for calcium dependent heterophilic binding to the desmocollins, whereas the intracellular domain is essential for binding to the desmosomal plaque protein, plakoglobin. Desmoglein 1 is synthesized exclusively in the suprabasal layers. Intact and functionally active desmoglein 1 is essential to epidermal integrity.

Product Info	
Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Western Blot (1-2ug/ml)

	NR	R	
250→			
150→			
100→			
75→			2ug loading NR=Non- reduced R=reduced
50→			
37→			
25→			
20→			
15→			
10→			

Fig. 1: SDS-PAGE Analysis Purified DSG1 Mouse Monoclonal Antibody (27B2). Confirmation of Integrity and Purity of Antibody.