

36-2239: Anti-ERCC1 / RAD10 (Tumor Progression Marker) Monoclonal Antibody(Clone: ERCC1/2318)

Clonality :	Monoclonal
Clone Name :	ERCC1/2318
Application :	ELISA, WB
Reactivity :	Human
Gene :	ERCC1
Gene ID :	2067
Uniprot ID :	P07992
Alternative Name :	COFS4; DNA excision repair protein ERCC1; ERCC1; Excision repair cross complementing 1; RAD10; UV20
Isotype :	Mouse IgG2b, kappa
Immunogen Information :	Recombinant fragment (around aa 191-281) of human ERCC1 protein (exact sequence is proprietary)

Description

Recognizes a protein of 110kDa, identified as Excision Repair Cross Complementing 1 (ERCC1). It is a mammalian nucleotide excision repair (NER) enzyme involved in repair of damaged DNA. ERCC1 is a homologous to RAD10 in *Saccharomyces cerevisiae*, which is required in mitotic intrachromosomal recombination and repair. ERCC1 is required in repair of cisplatin-induced DNA adducts and ultraviolet (UV)-induced DNA damage. High expression of ERCC1 has been linked to tumor progression in a variety of cancers including non-small cell lung cancer (NSCLC), squamous cell carcinoma of the head, ovarian cancer and esophageal cancer.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

ELISA (For coating, order antibody without BSA); Western Blot (1-2ug/ml);




Fig. 1: Formalin-fixed, paraffin-embedded human Pancreas stained with ERCC1 Mouse Monoclonal Antibody (ERCC1/2318).




Fig. 2: Formalin-fixed, paraffin-embedded human Pancreas stained with ERCC1 Mouse Monoclonal Antibody (ERCC1/2318).




Fig. 3: Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with ERCC1 Mouse Monoclonal Antibody (ERCC1/2318).




Fig. 4: SDS-PAGE Analysis Purified ERCC1 Mouse Monoclonal Antibody (ERCC1/2318). Confirmation of Purity and Integrity of Antibody.




Fig. 5: Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with ERCC1 Mouse Monoclonal Antibody (ERCC1/2318).




Fig. 6: Analysis of Protein Array containing more than 19,000 full-length human proteins using ERCC1 Mouse Monoclonal Antibody (ERCC1/2318). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.