

36-2277: Anti-Protocadherin FAT2 (FAT Atypical Cadherin 2) Monoclonal Antibody(Clone: 8C5)

Clonality :	Monoclonal
Clone Name :	8C5
Application :	FACS,IF,IHC
Reactivity :	Human
Gene :	FAT2
Gene ID :	2196
Uniprot ID :	Q9NYQ8
Alternative Name :	FAT tumor suppressor homolog 2 (HFAT2); cadherin family member 8 (CDHF8); cadherin-related family member 9 (CDHR9); multiple EGF-like domains protein 1; multiple epidermal growth factor-like domains 1 (MEGF1); multiple epidermal growth factor-like domains protein 1; protocadherin FAT2.
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Purified recombinant human FAT2 fusion protein.

Description

Recognizes a protein of 480kDa, which is identified as FAT2. The cadherins represent a family of Ca²⁺-dependent adhesion molecules that function to mediate cell-to-cell binding that is critical for the maintenance of structure and morphogenesis. Cadherins each contain a large extracellular domain at the N-terminus, which is characterized by a series of five homologous repeats, the most distal of which is thought to be responsible for binding specificity. The relatively short C-terminal intracellular domain interacts with a variety of cytoplasmic proteins, including -catenin, to regulate cadherin function. The cadherin superfamily includes cadherins, protocadherins, desmogleins and desmocollins. FAT2 (FAT tumor suppressor homolog 2) is a single-pass type I membrane protein that belongs to the protocadherin subfamily of cadherins. FAT2 contains one Laminin G-like domain, two EGF-like domains and 32 cadherin domains and is believed to function as a cell adhesion molecule, controlling cell proliferation and playing an important role in cerebellum development.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Cervical Carcinoma stained with Protocadherin FAT2 Monoclonal Antibody (8C5).




Fig. 2: Formalin-fixed, paraffin-embedded human Bladder Carcinoma stained with Protocadherin FAT2 Monoclonal Antibody (8C5).




Fig. 3: SDS-PAGE Analysis Purified Protocadherin FAT2 Monoclonal Antibody (8C5). Confirmation of Integrity and Purity of Antibody.