

36-2359: Anti-GAD1 / GAD67 (GABAergic Neuronal Marker) Monoclonal Antibody(Clone: GAD1/2391)

Clonality :	Monoclonal
Clone Name :	GAD1/2391
Application :	ELISA,WB,IHC
Reactivity :	Human
Gene :	GAD1
Gene ID :	2571
Uniprot ID :	Q99259
Alternative Name :	67kDa glutamic acid decarboxylase; CPSQ1; DCE1; GAD67; GAD1; Glutamate decarboxylase 1; SCP
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Recombinant human GAD1 (GAD67) protein fragment (around aa 72-135) (exact sequence is proprietary)

Description

This MAbs recognizes a protein of 67kDa, which is identified as glutamic acid decarboxylase 1 (GAD1). There are two forms of glutamic acid decarboxylases (GADs) that are found in the brain: GAD65 (also known as GAD2) and GAD67 (also known as GAD1). GAD65 and GAD67 are members of the group II decarboxylase family of proteins and are responsible for catalyzing the rate-limiting step in the production of GABA (-aminobutyric acid) from L-glutamic acid. Although both GADs are found in the brain, GAD65 localizes to synaptic vesicle membranes in nerve terminals, while GAD67 is distributed throughout the cell. GAD67 is responsible for the basal levels of GABA synthesis. In the case of a heightened demand for GABA in neurotransmission, GAD65 will transiently activate to assist in GABA production. The loss of GAD65 is detrimental and can impair GABA neurotransmission, however the loss of GAD67 is lethal.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

ELISA (For coating, order antibody without BSA); Western Blot (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT); (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Brain stained with GAD1 (GAD67) Mouse Monoclonal Antibody (GAD1/2391).




Fig. 2: SDS-PAGE Analysis Purified GAD1 (GAD67) Mouse Monoclonal Antibody (GAD1/2391). Confirmation of Integrity and Purity of Antibody.




Fig. 3: Western Blot Analysis of human Brain tissue lysate using GAD1 (GAD67) Mouse Monoclonal Antibody (GAD1/2391).




Fig. 4: Analysis of Protein Array containing more than 19,000 full-length human proteins using GAD1 (GAD67) Mouse Monoclonal Antibody (GAD1/2391) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.