

36-2383: Anti-Growth Hormone (Pituitary Marker) Monoclonal Antibody(Clone: GH/1371)

Clonality :	Monoclonal
Clone Name :	GH/1371
Application :	IHC
Reactivity :	Human
Gene :	GH1
Gene ID :	2688
Uniprot ID :	P01241
Alternative Name :	GH-N; GH1; Growth hormone 1; HG1; hGH-N; IGHD1B; Pituitary growth hormone; RNGHGP; Somatotropin
Isotype :	Mouse IgG2b, kappa
Immunogen Information :	A recombinant fragment (around aa58-187) of human Growth Hormone (GH) protein (exact sequence is proprietary)

Description

Pituitary growth hormone (GH) plays a crucial role in stimulating and controlling the growth, metabolism and differentiation of many mammalian cell types by modulating the synthesis of multiple mRNA species. These effects are mediated by the binding of GH to its membrane-bound receptor, GHR, and involve a phosphorylation cascade that results in the modulation of numerous signaling pathways. GH is synthesized by acidophilic or somatotrophic cells of the anterior pituitary gland. Anti-GH is a useful marker in classification of pituitary tumors and the study of pituitary disease (acromegaly).

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Pituitary stained with Growth Hormone Monoclonal Antibody (GH/1371).




Fig. 2: Formalin-fixed, paraffin-embedded human Pituitary stained with Growth Hormone Monoclonal Antibody (GH/1371).




Fig. 3: SDS-PAGE Analysis Purified Growth Hormone Mouse Monoclonal Antibody (GH/1371). Confirmation of Integrity and Purity of Antibody.




Fig. 4: Analysis of Protein Array containing more than 19,000 full-length human proteins using Growth Hormone Mouse Monoclonal Antibody (GH/1371). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.