

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-2420: Anti-CD13 / Aminopeptidase-N (Myeloid Cell Marker) Monoclonal Antibody(Clone: **APN/514)**

Clonality: Monoclonal **Clone Name:** APN/514 Application: **ELISA** Reactivity: Human Gene: ANPEP Gene ID: 290 **Uniprot ID:** P15144

Alanyl aminopeptidase; Aminopeptidase M; Aminopeptidase N; ANPEP; AP-M; APN; gp150;

Alternative Name: hAPN; Lap1; Microsomal aminopeptidase; Myeloid plasma membrane glycoprotein CD13;

p150; PEPN

Isotype: Mouse IgG1, kappa

Immunogen Information: Recombinant human CD13 protein

Description

This MAb recognizes an extracellular epitope of an integral membrane glycoprotein of 150kDa, identified as CD13. This antigen is present on most cells of myeloid origin including granulocytes, monocytes, mast cells, and GM-progenitor cells. It is also expressed by the majority of AML, CML in myeloid blast crisis, and in a smaller fraction of lymphoid leukemias. It is absent from normal lymphocytes, platelets and erythrocytes. CD13 is also present on fibroblasts; endothelial cells, epithelial cells from renal proximal tubules and intestinal brush border, bone marrow stromal cells, osteoclasts, and cells lining bile duct canaliculi. CD13 is identical to aminopeptidase N (APN), a prominent membrane-bound metalloprotease present on the surface of intestinal brush border and renal tubules. CD13 plays a role in metabolism of biologically active peptides, in phagocytosis, and in bactericidal/tumoricidal activities. It also serves as a receptor for human coronaviruses (HCV). The lineage-restricted pattern of expression of CD13 within the hemopoietic compartment sµggests that it may be important in myeloid cell differentiation.

Product Info

Amount: $20 \mu g / 100 \mu g$

200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS Content:

with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody Storage condition:

is stable for 24 months. Non-hazardous.

Application Note

ELISA (For coating, order antibody without BSA);