

36-2436: Anti-Glutathione S-Transferase Mu3 (GSTM3) Monoclonal Antibody(Clone: CPTC-GSTMu3-1)

Clonality :	Monoclonal
Clone Name :	CPTC-GSTMu3-1
Application :	IHC
Reactivity :	Human
Gene :	GSTM3
Gene ID :	2947
Uniprot ID :	P21266
Alternative Name :	brain GST antibody; brain type mu glutathione S transferase; GST class mu 3; GST5; GSTB; GSTM3-3; S (hydroxyalkyl)glutathione lyase M3; glutathione S transferase M3 (brain)
Isotype :	Mouse IgG2a, kappa
Immunogen Information :	Recombinant full-length human GSTM3 protein

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: SDS-PAGE Analysis Purified GST Mu3 Mouse Monoclonal Antibody (CPTC-GSTMu3-1). Confirmation of Purity and Integrity of Antibody.




Fig. 2: Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with Purified GST Mu3 Mouse Monoclonal Antibody (CPTC- GSTMu3-1).




Fig. 3: Analysis of Protein Array containing more than 19,000 full-length human proteins using Glutathione S-Transferase Mu3 (GSTM3) Mouse Monoclonal Antibody (CPTC-GSTMu3-1). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.