

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 36-2498: Anti-FOXA1 / HNF3A Monoclonal Antibody(Clone: FOXA1/1516)

Clonality: Monoclonal
Clone Name: FOXA1/1516

Application: IHC
Reactivity: Human
Gene: FOXA1
Gene ID: 3169
Uniprot ID: P55317

Alternative Name: Forkhead box protein A1 (FOXA1); Hepatocyte nuclear factor 3-alpha (HNF-3-alpha or HNF-3A

or HNF3A); Transcription factor 3A (TCF-3A or TCF3A)

**Isotype:** Mouse IgG2b, kappa

Immunogen Information: Recombinant human FOXA1 protein fragment (aa372-472) (exact sequence is proprietary)

## **Description**

The transcription factor Forkhead-box A1 (FOXA1), also known as hepatocyte nuclear factor 3-alpha, is a member of the FOX class of transcription factors.HNF-1 (), and HNF-6 compose, in part, a homeoprotein family designated the hepatocyte nuclear factor family. The various HNF-1 isoforms regulate transcription of genes in the liver as well as in other tissues such as kidney, small intestine and thymus. FOXA1 is expressed in normal breast ductal epithelium and other epithelium in different organs, such as lung, pancreas, bladder, prostate, and colon. Recently, FOXA1 has been shown to be a major determinant of estrogen-ER activity and endocrine response in breast cancer cells. FOXA1 expression correlates with estrogen receptor (ER)-positivity, especially in luminal subtype A breast cancers, which is associated with favorable prognosis. FOXA1 is useful in the sub-classification of breast carcinomas.

## **Product Info**

**Amount:** 20 μg / 100 μg

Content: 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS

with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

**Storage condition :** Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

## **Application Note**

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95&degC followed by cooling at RT for 20 minutes)




Fig. 1: Formalin-fixed, paraffin-embedded human Prostate stained with FOXA1 Mouse Monoclonal Antibody (FOXA1/1516).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig. 2: Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with FOXA1 Mouse Monoclonal Antibody (FOXA1/1516).




Fig. 3: Formalin-fixed, paraffin-embedded human Prostate (PIN) stained with FOXA1 Mouse Monoclonal Antibody (FOXA1/1516).




Fig. 4: Formalin-fixed, paraffin-embedded human Prostate stained with FOXA1 Mouse Monoclonal Antibody (FOXA1/1516).




Fig. 5: SDS-PAGE Analysis Purified FOXA1 Mouse Monoclonal Antibody (FOXA1/1516). Confirmation of Purity and Integrity of Antibody.