

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-2506: Anti-HSP60 (Heat Shock Protein 60) (Mitochondrial Marker) Monoclonal Antibody(Clone: LK2)

Clonality: Monoclonal

Clone Name: LK2

Application: IF,WB,IHC

Reactivity: Human, Mouse, Rat

 Gene :
 HSPD1

 Gene ID :
 3329

 Uniprot ID :
 P10809

60kDa chaperonin, 60kDa heat shock protein mitochondrial, Chaperonin, 60-KD (CPN60),

Alternative Name: GROEL, HLD4, HSP65, HSPD1, HuCHA60, Mitochondrial matrix protein P1, P60 lymphocyte

protein, Short heat shock protein 60 Hsp60s1, Spastic paraplegia 13 (SPG13)

Isotype: Mouse IgG1, kappa

Immunogen Information: Recombinant human HSP60 protein

Description

Recognizes a 60kDa protein, identified as the heat shock protein 60 (hsp60). Its epitope is localized between aa 383-419 of human hsp60. A wide variety of environmental and pathophysiological stressful conditions trigger the synthesis of a family of proteins known as heat shock proteins (hsp's), more appropriately called as stress response proteins (srp's). hsp60 is a potential antigen in a number of autoimmune diseases. In human arthritis and in experimentally induced arthritis in animals, disease development coincides with the development of immune reactivity directed against not only bacterial hsp60, but also against its mammalian homolog.

Product Info

Amount : 20 μg / 100 μg

Content: 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS

with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml.

Storage condition : Antibody with azide - store at 2 to 8°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Immunofluorescence (1-2ug/ml); Western Blot (1-2ug/ml); ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),




Fig. 1: Immunofluorescent staining of MEOH-fixed HeLa cells using HSP60 Mouse Monoclonal Antibody (LK2) followed by goat anti-Mouse labeled with CF488 (Green); Nuclei are labeled with NucSpot red).