

### 36-2520: Anti-CD50 / ICAM3 Monoclonal Antibody(Clone: ICAM3/2873R)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	ICAM3/2873R
<b>Application :</b>	IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	ICAM3
<b>Gene ID :</b>	3385
<b>Uniprot ID :</b>	P32942
<b>Alternative Name :</b>	CD50; CDW 50; CDw50; ICAM 3; ICAM-3; ICAM-R; ICAM3; ICAM3_HUMAN; ICAMR; Intercellular adhesion molecule 3
<b>Isotype :</b>	Rabbit IgG
<b>Immunogen Information :</b>	Recombinant full-length human ICAM3 protein

#### Description

Recognizes the D1 domain of an N-glycosylated glycoprotein of 120kDa with intra-chain disulfide bonds, identified as CD50 or ICAM-3 (Leucocyte Workshop VI). CD50 is the major ligand for LFA-1 (CD11a/CD18) and may have signaling role to increase adhesion. It is expressed on thymocytes and T lymphocytes and is resistant to treatment with phosphatidylinositol (PI) phospholipase C. This MAbs is excellent for staining of formalin/paraffin tissues.

#### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

#### Application Note

Blocks binding of CD11a (LFA-1) to CD50 (ICAM-3) (Order Ab without Azide); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);




Fig. 1: SDS-PAGE Analysis Purified CD50 Rabbit Recombinant Monoclonal Antibody (ICAM3/2873R). Confirmation of Purity and Integrity of Antibody.




Fig. 2: Formalin-fixed, paraffin-embedded human Tonsil stained with CD50 Rabbit Recombinant Monoclonal Antibody (ICAM3/2873R).