9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

36-2795: Anti-MMP9 (Matrix Metalloproteinase 9) Monoclonal Antibody(Clone: rMMP9/1769)

Clonality :	Monoclonal
Clone Name :	rMMP9/1769
Application :	IHC
Reactivity :	Human
Gene :	MMP9
Gene ID :	4318
Uniprot ID :	P14780
Alternative Name :	82 kDa matrix metalloproteinase-9; 92 kDa gelatinase; 92 kDa type IV collagenase; CLG 4B; CLG4B; Collagenase Type 4 beta; Collagenase type IV 92 KD; EC 3.4.24.35; Gelatinase 92 KD; Gelatinase B; Gelatinase beta; GelatinaseB; GELB; Macrophage gelatinase; MANDP2; Matrix metallopeptidase 9 (gelatinase B, 92kDa gelatinase, 92kDa type IV collagenase); Matrix Metalloproteinase 9; MMP 9; MMP9; Type V collagenase
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Recombinant full-length human MMP-9 protein

Description

The matrix metalloproteinases (MMP) are a family of peptidase enzymes responsible for the degradation of extracellular matrix components, including collagen, gelatin, fibronectin, laminin and proteoglycan. Transcription of MMP genes is differentially activated by phorbol ester, lipopolysaccharide (LPS) or staphylococcal enterotoxin B (SEB). MMP catalysis requires both calcium and zinc. MMP-9 (also designated 92 kDa type IV collagenase or gelatinase B) has been shown to degrade bone collagens in concert with MMP-1 (also designated interstitial collagenase, fibroblast collagenase or collagenase-1), and cysteine proteases and may play a role in bone osteoclastic resorption. MMP-1 is down-regulated by p53, and abnormality of p53 expression may contribute to joint degradation in rheumatoid arthritis by regulating MMP-1 expression.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);

For Research Use Only. Not for use in diagnostic/therapeutics procedures.

9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com




Fig. 1: Formalin-fixed, paraffin-embedded human spleen stained with MMP9 Mouse Recombinant Monoclonal Antibody (rMMP9/1769).




Fig. 2: SDS-PAGE Analysis Purified MMP9 Mouse Recombinant Monoclonal Antibody (rMMP9/1769). Confirmation of Integrity of Purity of Antibody.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.