

## 36-2805: Anti-MTAP (Tumor Suppressor Marker) Monoclonal Antibody(Clone: rMTAP/1813)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	rMTAP/1813
<b>Application :</b>	IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	MTAP
<b>Gene ID :</b>	4507
<b>Uniprot ID :</b>	Q13126
<b>Alternative Name :</b>	BDMF; DMSFH; DMSMFH; Epididymis luminal protein 249; HEL249; LGMBF; MeSAdo phosphorylase; Methylthioadenosine phosphorylase; MSAP; MTA phosphorylase; MTAPase; S-methyl-5''-thioadenosine phosphorylase
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Recombinant human MTAP protein fragment (aa97-196) (exact sequence is proprietary)

### Description

Recognizes a protein of 31kDa, which is identified as MTAP (5'-deoxy-5'-methylthioadenosine phosphorylase). It catalyzes the reversible phosphorolysis of methylthioadenosine, which is important in polyamine metabolism and for the salvage of adenine and methionine. The gene encoding MTAP is linked to the tumor suppressor gene, p16INK4A. Deficient levels of MTAP can occur in cancers primarily through co-deletion of the MTAP gene and the p16INK4A gene. Cells expressing MTAP and possessing adenine salvage pathway activity may be less susceptible to malignancy due to growth-inhibitory actions of agents (e.g. antifolates), whose mechanism of action, in part, involves this de novo purine pathway.

### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

### Application Note

Immunohistochemistry (Formalin-fixed) (1-2µg/ml for 30 minutes at RT),(Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95&degC followed by cooling at RT for 20 minutes),




Fig. 1: Formalin-fixed, paraffin-embedded human Kidney stained with MTAP Recombinant Mouse Monoclonal Antibody (rMTAP/1813).




Fig. 2: Formalin-fixed, paraffin-embedded human Kidney stained with MTAP Recombinant Mouse Monoclonal Antibody (rMTAP/1813).




Fig. 3: SDS-PAGE Analysis Purified MTAP Recombinant Mouse Monoclonal Antibody (rMTAP/1813). Confirmation of Purity and Integrity of Antibody.