

36-2821: Anti-MUC3 (Mucin 3) Monoclonal Antibody(Clone: MUC3/2992R)

Clonality :	Monoclonal
Clone Name :	MUC3/2992R
Application :	IHC
Reactivity :	Human
Gene :	MUC3A
Gene ID :	4584; 57876
Uniprot ID :	Q02505; Q9H195
Alternative Name :	Intestinal mucin-3A; intestinal mucin-like; MUC3; MUC3A; MUC3B; mucin 3, intestinal; mucin 3A, cell surface associated; mucin 3A, intestinal; mucin 3B, cell surface associated
Isotype :	Rabbit IgG
Immunogen Information :	A synthetic peptide of 35 amino acids, SIB35 (C-HSTPSFTSSITTTETTSHTSPSTSSITTTETTS), which contains two of the MUC3 tandem repeats, coupled to KLH.

Description

It recognizes a protein of HMW, identified as mucin 3 glycoprotein (MUC3). Its epitope localizes between aa SITTE. This MAb shows no cross-reaction with human milk fat globule membranes, MUC1, or MUC2. MUC3 is distributed in colon and rectum, and is also present to a lesser extent in breast, lung and salivary gland tissues. The Mucins are a family of highly glycosylated, secreted proteins with a basic structure consisting of a variable number of tandem repeats (VNTRs) encoded by 60 base pairs (Mucin 1), 69 base pairs (Mucin 2) and 51 base pairs (Mucin 3). The number of repeats is highly polymorphic and varies among different alleles. Mucin 1 proteins are expressed as type I membrane proteins in addition to secreted forms. Mucin 1 is aberrantly expressed in epithelial tumors including breast carcinomas. Mucin 2 coats the epithelia of the intestines and airways and is associated with colonic tumors. Mucin 3 is a major component of various mucus gels and is broadly expressed in normal and tumor cells.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2µg/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes)




Fig. 1: Formalin-fixed, paraffin-embedded human Gastric Carcinoma stained with MUC3 Rabbit Recombinant Monoclonal Antibody (MUC3/2992R).




Fig. 2: SDS-PAGE Analysis Purified MUC3 Rabbit Recombinant Monoclonal Antibody (MUC3/2992R). Confirmation of Purity and Integrity of Antibody.