9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

36-2831: Anti-c-Myc Oncoprotein Monoclonal Antibody(Clone: MYC699)

Clonality :	Monoclonal
Clone Name :	MYC699
Application :	ELISA,FACS,IF
Reactivity :	Human
Gene :	MYC
Gene ID :	4609
Uniprot ID :	P01106
Alternative Name :	Class E basic helix-loop-helix protein 39 (bHLHe39), MRTL, Myc2, Niard, Nird, Proto-oncogene c-Myc, RNCMYC, Transcription factor p64, Transcriptional regulator Myc-A, V-Myc avian myelocytomatosis viral oncogene homolog
Isotype :	Mouse IgG1, kappa
Immunogen Information :	A synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQL-RNSCA) from C-terminus of human c-myc, coupled to KLH

Description

The c-Myc protein is a transcription factor, which is encoded by the c-Myc gene on human chromosome 8q24. c-Myc is commonly activated in a variety of tumor cells and plays an important role in cellular proliferation, differentiation, apoptosis and cell cycle progression. The phosphorylation of c-Myc has been investigated and previous studies have sÂµggested a functional association between phosphorylation at Thr58/Ser62 by glycogen synthase kinase 3, cyclin dependent kinase, ERK2 and C-Jun N terminal Kinase (JNK) in cell proliferation and cell cycle regulation. Studies also have shown that c-Myc is essential for tumor cell development in vasculogenesis and angiogenesis that distribute blood throÂµghout the cells, and which broÂµght extensive attention in the development of new therapeutic approach for cancer treatment.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

ELISA (Use Ab at 2-4ug/ml for coating) (Order Ab without BSA); Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml);



Fig. 1: SDS-PAGE Analysis Purified MYC Mouse Monoclonal Antibody (MYC699). Confirmation of Integrity and Purity of Antibody.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.