

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-2833: Anti-c-Myc Oncoprotein Monoclonal Antibody(Clone: MYC2895R)

Clonality: Monoclonal MYC2895R Clone Name: Application: ELISA, FACS, IF

Reactivity: Human Gene: **MYC** Gene ID: 4609 **Uniprot ID:** P01106

Class E basic helix-loop-helix protein 39 (bHLHe39), MRTL, Myc2, Niard, Nird, Proto-oncogene c-

Myc, RNCMYC, Transcription factor p64, Transcriptional regulator Myc-A, V-Myc avian **Alternative Name:**

myelocytomatosis viral oncogene homolog

Rabbit IgG Isotype:

A synthetic peptide, corresponding to aa 408-439

Immunogen Information: (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-myc, coupled to

Description

It recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This MAb shows no cross-reaction with v-myc. cmyc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over-expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.

Product Info

20 μg / 100 μg Amount:

200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with Content:

0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is Storage condition:

stable for 24 months. Non-hazardous.

Application Note

ELISA (For coating, order antibody without BSA); Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml);




Fig. 1: SDS-PAGE Analysis Purified c-Myc Rabbit Recombinant Monoclonal Antibody (MYC2895R). Confirmation of Purity and Integrity of Antibody.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig. 2: Flow Cytometric Analysis of K562 cells using c-Myc Rabbit Recombinant Monoclonal Antibody (MYC2895R) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype control (Red).