

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 36-2849: Anti-NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) Monoclonal Antibody(Clone: NGFR/2550R)

Clonality: Monoclonal
Clone Name: NGFR/2550R
Application: IHC,WB
Reactivity: Human
Gene: NGFR
Gene ID: 4804
Uniprot ID: P08138

CD271; Gp80-LNGFR; Low affinity nerve growth factor receptor; Low affinity neurotrophin

**Alternative Name:** receptor p75NTR; Nerve growth factor receptor (NGFR); p75 ICD; p75 Neurotrophin receptor;

Tumor necrosis factor receptor superfamily member 16 (TNFRSF16)

**Isotype:** Rabbit IgG

Immunogen Information: Recombinant human p75 NGFR protein fragment (around aa 281-421) (exact sequence is

proprietary)

## **Description**

It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts.

## **Product Info**

**Amount :**  $20 \mu g / 100 \mu g$ 

Content: 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS

with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

**Storage condition :** Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody

is stable for 24 months. Non-hazardous.

## **Application Note**

Western Blot (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig. 1: Formalin-fixed, paraffin-embedded human Melanoma stained with NGFR Rabbit Recombinant Monoclonal Antibody (NGFR/2550R).




Fig. 2: SDS-PAGE Analysis Purified NGFR Rabbit Recombinant Monoclonal Antibody (NGFR/2550R). Confirmation of Integrity and Purity of Antibody.