9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

36-2900: Anti-PAX5 / BSAP (Early B-Cell Marker) Monoclonal Antibody(Clone: PAX5/3735)

Clonality :	Monoclonal
Clone Name :	PAX5/3735
Application :	IHC
Reactivity :	Human
Gene :	PAX5
Gene ID :	5079
Uniprot ID :	Q02548
Alternative Name :	B cell lineage specific activator; BSAP; EBB-1; KLP; Paired box 5
Isotype :	Mouse IgG1, lambda
Immunogen Information :	A recombinant fragment (around aa 235-382) of human PAX5 protein (exact sequence is proprietary)

Description

The specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 21,000, full-length human proteins. The PAX5 gene is a member of thepaired box (PAX)family of transcription factors. The central feature of this gene family is a novel, highly conservedDNA-binding domain, known as thepaired box. The PAX proteins are important regulators in early development, and alterations in the expression of their genes are thoÂµght to contribute toneoplastic transformation. The PAX5 gene encodes theB-celllineage specific activator protein (BSAP) that is expressed at early, but not late stages of B-cell differentiation. Its expression has also been detected in developing CNS and testis; therefore, PAX5 gene product may not only play an important role in B-cell differentiation, but also in neural development andspermatogenesis.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);



Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil stained with PAX5 Mouse Monoclonal Antibody (PAX5/3735).

For Research Use Only. Not for use in diagnostic/therapeutics procedures.

9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com




Fig. 2: Analysis of Protein Array containing more than 21,000 full-length human proteins using PAX5 Mouse Monoclonal Antibody (PAX5/3735) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.