

36-2921: Anti-CD31 / PECAM-1 (Endothelial Cell Marker) Monoclonal Antibody(Clone: 158-2B3)

Clonality :	Monoclonal
Clone Name :	158-2B3
Application :	FACS,IF
Reactivity :	Human
Gene :	PECAM1
Gene ID :	5175
Uniprot ID :	P16284
Alternative Name :	EndoCAM; PECA1; Platelet Endothelial Cell Adhesion Molecule 1; GPIIA'
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Stimulated human leukocytes (Workshop VI)

Description

CD31 (PECAM-1) is a transmembrane glycoprotein member of the immunoglobulin supergene family of adhesion molecules. CD31 is expressed by stem cells of the hematopoietic system and is primarily used to identify and concentrate these cells for experimental studies as well as for bone marrow transplantation. Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 MAb reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml);




Fig. 1: Flow Cytometric Analysis of paraformaldehyde-fixed Jurkat cells using CD31 Mouse Monoclonal Antibody (158-2B3) followed by goat anti- Mouse- IgG-CF488 (Blue); Isotype Control (Red).




Fig. 2: SDS-PAGE Analysis Purified CD31 Mouse Monoclonal Antibody (158-2B3). Confirmation of Integrity and Purity of Antibody.




Fig. 3: Immunofluorescent staining of PFA-fixed Jurkat cells using CD31 Mouse Monoclonal Antibody (158-2B3) followed by goat anti-Mouse IgG conjugated with CF488 (green). Nuclei are labeled with RedDot (red).