

36-2987: Anti-DOG-1 / TMEM16A (Gastrointestinal Stromal Tumor Marker) Monoclonal Antibody(Clone: DG1/1484)

Clonality :	Monoclonal
Clone Name :	DG1/1484
Application :	IHC
Reactivity :	Human
Gene :	TMEM16A
Gene ID :	55107
Uniprot ID :	Q5XX6
Alternative Name :	Anoctamin 1; Calcium Activated Chloride Channel; Discovered On Gastrointestinal Stromal Tumors Protein 1; TAOS2; ORAOV2; TMEM16A
lsotype :	Mouse IgG2b, kappa
Immunogen Information :	Recombinant human DOG-1 protein fragment (aa 2-101) (exact sequence is proprietary)

Description

Expression of DOG-1 protein is elevated in the gastrointestinal stromal tumors (GIST's), c-kit signaling-driven mesenchymal tumors of the GI tract. DOG-1 is rarely expressed in other soft tissue tumors, which, due to appearance, may be difficult to diagnose. Immunoreactivity for DOG-1 has been reported in 97.8 percent of scorable GIST's, including all c-kit negative GIST's. Overexpression of DOG-1 has been sµggested to aid in the identification of GISTs, including Platelet-Derived Growth Factor Receptor Alpha mutants that fail to express c-kit antigen. The overall sensitivity of DOG1 and c-kit in GIST's is nearly identical: 94.4% vs. 94.7%.

Product Info

Amount :	20 μg / 100 μg
Content :	200 μg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);



Fig. 1: Formalin-fixed, paraffin-embedded human GIST stained with DOG-1 Mouse Monoclonal Antibody (DG1/1484).

∗ abeomics

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com

kDa	R	NR	
250 —			
150 —		-	
100 —			
75	-		2ug loading
50	-	•	NR=Non-
37			R=reduced
25 —		-	
20			
15 —			
10			

Fig. 2: SDS-PAGE Analysis Purified DOG-1 Mouse Monoclonal Antibody (DG1/1484). Confirmation of Integrity and Purity of Antibody



Fig. 3: Analysis of Protein Array containing more than 19,000 full-length human proteins using DOG-1 Mouse Monoclonal Antibody (DG1/1484). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.