

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-3064: Anti-CD45 / LCA (B-Cell Marker) Monoclonal Antibody(Clone: PTPRC/1461)

Clonality: Monoclonal
Clone Name: PTPRC/1461
Application: FACS,IF,IHC
Reactivity: Human
Gene: PTPRC
Gene ID: 5788
Uniprot ID: P08575

B220, CD45R, GP180, Leukocyte common antigen (LCA), Loc, Ly-5, Lyt-4, Protein tyrosine

Alternative Name: phosphatase receptor type C (PTPRC), Receptor-type tyrosine-protein phosphatase C, T200

glycoprotein

Isotype: Mouse IgG2a, kappa **Immunogen Information**: Human B-lymphocytes

Description

CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.

Product Info

Amount : 20 μg / 100 μg

Content: 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with

0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage condition:

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is

stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil stained with CD45 Monoclonal Antibody (PTPRC/1461).



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Fig. 2: Immunofluorescence Analysis of PFA-fixed Jurkat cells labeling CD45RA with CD45 Monoclonal Antibody (PTPRC/1461) followed by Goat anti-Mouse IgG-CF488 (Green). The nuclear counterstain is Redot.




Fig. 3: Flow Cytometric Analysis of PFA-fixed Jurkat cells using CD45 Monoclonal Antibody (PTPRC/1461) followed by Goat anti- Mouse- IgG-CF488 (Blue); Isotype Control (Red).




Fig. 4: SDS-PAGE Analysis Purified CD45 Monoclonal Antibody (PTPRC/1461). Confirmation of Integrity and Purity of Antibody