

### 36-3119: Anti-S100A4 / Metastasin / Calvasculin (Marker of Tumor Metastasis) Monoclonal Antibody(Clone: S100A4/1481)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	S100A4/1481
<b>Application :</b>	IHC,FACS,WB,IF
<b>Reactivity :</b>	Human
<b>Gene :</b>	S100A4
<b>Gene ID :</b>	6275
<b>Uniprot ID :</b>	P26447
<b>Alternative Name :</b>	S100A4; S100 calcium-binding protein A4; Calvasculin; CAPL; Fibroblast specific protein 1 (FSP1); Leukemia multidrug resistance associated protein; Malignant transformation suppression 1 (MTS1); Metastasin; Placental calcium-binding protein
<b>Isotype :</b>	Mouse IgG1, kappa
<b>Immunogen Information :</b>	Recombinant fragment (around aa 1-200) of human S100A4 protein (exact sequence is proprietary)

#### Description

S100A4 belongs to the S100 super-family of proteins containing 2 EF-hand calcium-binding domains. S100 genes include at least 25 members, including S100A1-S100A18, trichohyalin, filaggrin, repetin, S100P, and S100Z. S100A4 exerts its function via direct interaction with a number of proteins including P53, P63, non-muscle myosin IIA,  $\alpha$ 6 integrin, and liprin b1. S100A4 is overexpressed in highly metastatic cancers, which makes it useful as a marker of tumor progression.

#### Product Info

<b>Amount :</b>	20 $\mu$ g / 100 $\mu$ g
<b>Content :</b>	200 $\mu$ g/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

#### Application Note

Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Western Blot (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Placenta stained with S100A4 Mouse Monoclonal Antibody (S100A4/1481).




Fig. 2: Immunofluorescence staining of HeLa cells using S100A4 Mouse Monoclonal Antibody (S100A4/1481) followed by goat anti-Mouse IgG conjugated to CF488 (green). Nuclei are stained with Reddot.




Fig. 3: Flow Cytometric Analysis of T98G cells using S100A4 Mouse Monoclonal Antibody (S100A4/1481) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).




Fig. 4: Flow Cytometric Analysis of A549 cells using S100A4 Mouse Monoclonal Antibody (S100A4/1481) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).




Fig. 5: Western Blot of HeLa, A549 and A375 cell lysates using S100A4 Mouse Monoclonal Antibody (S100A4/1481).




Fig. 6: Western Blot Analysis of HeLa cell lysate using S100A4 Mouse Monoclonal Antibody (S100A4/1481).




Fig. 7: SDS-PAGE Analysis Purified S100A4 Mouse Monoclonal Antibody (S100A4/1481). Confirmation of Integrity and Purity of Antibody