

36-3120: Anti-S100A4 / Metastasin / Calvasculin (Marker of Tumor Metastasis) Monoclonal Antibody(Clone: S100A4/1482)

Clonality :	Monoclonal
Clone Name :	S100A4/1482
Application :	IHC,FACS,WB,IF
Reactivity :	Human
Gene :	S100A4
Gene ID :	6275
Uniprot ID :	P26447
Alternative Name :	S100A4; S100 calcium-binding protein A4; Calvasculin; CAPL; Fibroblast specific protein 1 (FSP1); Leukemia multidr�ug resistance associated protein; Malignant transformation suppression 1 (MTS1); Metastasin; Placental calcium-binding protein
Isotype :	Mouse IgG1, kappa
Immunogen Information :	Recombinant fragment of human S100A4 protein (aa1-200) (exact sequence is proprietary)

Description

S100A4 belongs to the S100 super-family of proteins containing 2 EF-hand calcium-binding domains. S100 genes include at least 25 members, including S100A1-S100A18, trichohyalin, filaggrin, repetin, S100P, and S100Z. S100A4 exerts its function via direct interaction with a number of proteins including P53, P63, non-muscle myosin IIA,  4 integrin, and liprin b1. S100A4 is overexpressed in highly metastatic cancers, which makes it useful as a marker of tumor progression.

Product Info

Amount :	20 �g / 100 �g
Content :	200 �g/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8�C. Antibody without azide - store at -20 to -80�C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry (1-2 g/million cells); Immunofluorescence (1-2 g/ml); ,Western Blot (1-2 g/ml); ,Immunohistochemistry (Formalin-fixed) (1-2 g/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Placenta stained with S100A4 Mouse Monoclonal Antibody (S100A4/1482).




Fig. 2: Western Blot of HeLa, A549 and A375 cell lysate using S100A4 Mouse Monoclonal Antibody (S100A4/1482).




Fig. 3: SDS-PAGE Analysis Purified S100A4 Mouse Monoclonal Antibody (S100A4/1482). Confirmation of Integrity and Purity of Antibody.




Fig. 4: Western Blot Analysis of HeLa cell lysate using S100A4 Mouse Monoclonal Antibody (S100A4/1482).




Fig. 5: Flow Cytometric Analysis of T98G cells using S100A4 Mouse Monoclonal Antibody (S100A4/1482) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).




Fig. 6: Flow Cytometric Analysis of A549 cells using S100A4 Mouse Monoclonal Antibody (S100A4/1482) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).