

36-3162: Anti-GLUT-1 (Tumor Progression and Mesothelioma Marker) Monoclonal Antibody(Clone: GLUT1/3132R)

Clonality :	Monoclonal
Clone Name :	GLUT1/3132R
Application :	ELISA,FACS,IHC
Reactivity :	Human
Gene :	SLC2A1
Gene ID :	6513
Uniprot ID :	P11166
Alternative Name :	Erythrocyte/hepatoma glucose transporter; Glucose transporter type-1; GLUT1; GLUT1DS; GLUTB; GT1; GTG1; Gtg3; HepG2 glucose transporter; PED; RATGTG1; Solute carrier family 2, facilitated glucose transporter member 1 (SLC2A1)
Isotype :	Rabbit IgG
Immunogen Information :	Recombinant fragment of human GLUT1 protein (around aa 203-305) (exact sequence is proprietary)

Description

Recognizes a protein of 55kDa, which is identified as GLUT-1. Glucose transporters are integral membrane glycoproteins involved in transporting glucose into most cells. There are many types of glucose transport carrier proteins, designated as Glut-1 to Glut-12. Glut-1 is a major glucose transporter in the mammalian blood-brain barrier. It is expressed in high density on the membranes of human erythrocytes and the brain capillaries that comprise the blood-brain barrier. Glut-1 is expressed at variable levels in many human tissues. Overexpression of Glut-1 has been linked to tumor progression or poor survival of patients with carcinomas of the colon, breast, cervical, lung, bladder and mesothelioma. Glut-1 is a sensitive and specific marker for the differentiation of malignant mesothelioma (positive) from reactive mesothelium (negative).

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

ELISA (For coating use Ab at 1-2ug/ml order Ab without BSA); Flow Cytometry (1-2ug/million cells); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Tongue stained with GLUT-1 Recombinant Rabbit Monoclonal Antibody (GLUT1/3132R).




Fig. 2: SDS-PAGE Analysis Purified GLUT-1 Recombinant Rabbit Monoclonal Antibody (GLUT1/3132R). Confirmation of Purity and Integrity of Antibody.




Fig. 3: Flow Cytometric Analysis of K562 cells using GLUT-1 Recombinant Rabbit Monoclonal Antibody (GLUT1/3132R) followed by goat anti-rabbit IgG-CF488 (Blue); Isotype Control (Red).