

### 36-3192: Anti-Spectrin Alpha 1 (Erythrocyte Marker) Monoclonal Antibody(Clone: SPTA1/1810)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	SPTA1/1810
<b>Application :</b>	ELISA,IHC
<b>Reactivity :</b>	Human
<b>Gene :</b>	SPTA1
<b>Gene ID :</b>	6708
<b>Uniprot ID :</b>	P02549
<b>Alternative Name :</b>	Alpha I spectrin; Elliptocytosis 2 (EL2); Erythrocyte alpha spectrin; Erythroid alpha-spectrin; HPP; HS3; Spectrin alpha erythrocytic 1; SPH3; SPTA1
<b>Isotype :</b>	Mouse IgG2c, kappa
<b>Immunogen Information :</b>	Recombinant human SPTA fragment (around aa356-475) (exact sequence is proprietary)

#### Description

Spectrin is an actin crosslinking and molecular scaffold protein that links the plasma membrane to the actin cytoskeleton, and functions in the determination of cell shape, arrangement of transmembrane proteins, and organization of organelles. Spectrins function as membrane organizers and stabilizers, composed of non-homologous ) are present in other somatic cells. The spectrin tetramers in erythrocytes act as barriers to lateral diffusion, but spectrin dimers seem to lack this function.

#### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

#### Application Note

ELISA (Use Ab at 2-4ug/ml for coating) (Order Ab without BSA); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Placenta stained with Spectrin alpha 1 (SPTA1) Mouse Monoclonal Antibody (SPTA1/1810).




Fig. 2: Formalin-fixed, paraffin-embedded human Tonsil stained with Spectrin alpha 1 (SPTA1) Mouse Monoclonal Antibody (SPTA1/1810).




Fig. 3: SDS-PAGE Analysis of Purified Spectrin alpha 1 Mouse Monoclonal Antibody (SPTA1/1810). Confirmation of Purity and Integrity.




Fig. 4: Analysis of Protein Array containing more than 19,000 full-length human proteins using Spectrin, alpha 1 Mouse Monoclonal Antibody (SPTA1/1810) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.