

36-3362: Anti-Ezrin / p81 Monoclonal Antibody(Clone: SPM244)

Clonality :	Monoclonal
Clone Name :	SPM244
Application :	WB,IHC
Reactivity :	Human
Gene :	Ezrin
Gene ID :	7430
Uniprot ID :	P15311
Alternative Name :	Villin-2; CVIL; Epididymis secretory protein Li 105; EZR; p81; VIL2; Cytovillin; Cytovillin 2; HEL S 105; DKFZp762H157
Isotype :	Mouse IgG2b, kappa
Immunogen Information :	Recombinant human full-length protein

Description

Ezrin, Moesin and Radixin belong to a family of highly homologous Actinassociated proteins that are localized just beneath the plasma membrane. The proteins are believed to be involved in the mediation of interactions between cytoskeletal and membrane proteins. Ezrin serves as a major cytoplasmic substrate of various protein-tyrosine kinases, including the epidermal growth factor receptor. Ezrin has also been identified as a cAMP-dependent protein kinase (A-kinase) anchoring protein and designated AKAP78. Moesin and Radixin share over 70% homology with Ezrin and are coexpressed within various cell types. Despite the high degree of homology, the three proteins exhibit a distinct receptor-specific pattern of phosphorylation. Overexpression of Ezrin predicts the poor prognosis of gastric adenocarcinoma.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Western Blot (1-2ug/ml); Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes);




Fig. 1: Formalin-fixed, paraffin-embedded human Placenta stained with Ezrin Mouse Monoclonal Antibody (SPM244).