9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

36-3460: Anti-CD2 / Lymphocyte Function Antigen 2 (LFA-2) Monoclonal Antibody(Clone: 1E7E8.G4)

Clonality :	Monoclonal
Clone Name :	1E7E8.G4
Application :	Functional Assay,FACS,IF,WB
Reactivity :	Human
Gene :	CD2
Gene ID :	914
Uniprot ID :	P06729
Alternative Name :	Erythrocyte receptor; LFA-2; LFA-3 receptor; Ly-37; Lymphocyte function antigen 2; Rosette receptor; SRBC; T-cell surface antigen CD2; T-cell surface antigen T11/Leu-5; T11
Isotype :	Mouse IgG2a, kappa
Immunogen Information :	Human Thymocytes/Sezary T cells

Description

CD2 interacts throÂµgh its amino-terminal domain with the extracellular domain of CD58 (also designated CD2 ligand) to mediate cell adhesion. CD2/CD58 binding can enhance antigen-specific T cell activation. CD2 is a transmembrane glycoprotein that is expressed on peripheral blood T lymphocytes, NK cells and thymocytes. CD58 is a heavily glycosylated protein with a broad tissue distribution in hematopoietic and other cells, including endothelium. Interaction between CD2 and its counter receptor LFA3 (CD58) on opposing cells optimizes immune system recognition, thereby facilitating communication between helper T lymphocytes and antigen-presenting cells, as well as between cytolytic effectors and target cells.

Product Info

Amount :	20 µg / 100 µg
Content :	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage condition :	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Functional Studies (Order Ab without Azide);,Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); ,Western Blot (1-2ug/ml);,




Fig. 1: SDS-PAGE Analysis Purified CD2 Mouse Monoclonal Antibody (1E7E8.G4). Confirmation of Integrity and Purity of Antibody.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.

9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com




Fig. 2: Western Blot Analysis of Jurkat cell lysate using CD2 Mouse Monoclonal Antibody (1E7E8.G4).




Fig. 3: Immunofluorescence staining of MOLT-4 cells using CD2 Mouse Monoclonal Antibody (1E7E8.G4) followed by goat anti-Mouse IgG conjµgated to CF488 (green). Nuclei are stained with Reddot.




Fig. 4: Flow Cytometric Analysis of MOLT4 cells using CD2 Mouse Monoclonal Antibody (1E7E8.G4) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).

For Research Use Only. Not for use in diagnostic/therapeutics procedures.