

### 36-3534: Anti-CD19 (B-Lymphocyte Marker) Monoclonal Antibody(Clone: PDR134)

<b>Clonality :</b>	Monoclonal
<b>Clone Name :</b>	PDR134
<b>Application :</b>	ELISA,FACS,IF,WB
<b>Reactivity :</b>	Human
<b>Gene :</b>	CD19
<b>Gene ID :</b>	930
<b>Uniprot ID :</b>	P15391
<b>Alternative Name :</b>	B-lymphocyte antigen CD19; B-lymphocyte surface antigen B4; CVID3; Leu-12; Lymphocyte Surface Antigen; T-cell surface antigen Leu-12
<b>Isotype :</b>	Mouse IgM, kappa
<b>Immunogen Information :</b>	Pokeweed-stimulated Daudi and Raji cells

#### Description

CD19 is a transmembrane glycoprotein that contains two extracellular immunoglobulin-like domains. CD19 is present in both benign and malignant B-cells and is considered to be the most reliable surface marker of this lineage over a wide range of maturational stages. In normal lymphoid tissue, CD19 is observed in germinal centers, in mantle zone cells, and in scattered cells of the inter-follicular areas. Anti-CD19 exhibits an overall immunoreactivity pattern similar to those of the antibodies against CD20 and CD22. However, in contrast to CD20, expression of CD19 is continuous throughout B-cell development and throughout terminal differentiation of B-cells into plasma cells. Anti-CD19 positivity is seen in the vast majority of B-cell neoplasms commonly at a lower intensity than normal B-cell counterparts. Plasma cell neoplasms are nearly always negative, as are T-cell neoplasms.

#### Product Info

<b>Amount :</b>	20 µg / 100 µg
<b>Content :</b>	200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
<b>Storage condition :</b>	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous.

#### Application Note

ELISA (For coating, order antibody without BSA); Flow Cytometry (1-2ug/million cells); Immunofluorescence (1-2ug/ml); Western Blot (1-2ug/ml);