

36-3543: Anti-CD20 / MS4A1 (B-Cell Marker) Monoclonal Antibody(Clone: L26)

| | |
|--------------------------------|--|
| Clonality : | Monoclonal |
| Clone Name : | L26 |
| Application : | FACS,IF,WB,IHC |
| Reactivity : | Human |
| Gene : | MS4A1 |
| Gene ID : | 931 |
| Uniprot ID : | P11836 |
| Alternative Name : | APY; ATOPY; B-lymphocyte cell-surface antigen B1; Bp35; Fc epsilon receptor I beta chain; Fc Fragment of IgE high affinity I receptor for beta polypeptide; FCER1B; High affinity immunoglobulin epsilon receptor subunit beta; IgE Fc receptor subunit beta; IGEL; IGER; IGHF; Leukocyte surface antigen Leu-16; Ly44; MS4A1; MS4A2 |
| Isotype : | Mouse IgG2a, kappa |
| Immunogen Information : | Human tonsil B cells |

Description

Recognizes a protein of 30-33kDa, which is identified as CD20. Its epitope is located in the cytoplasmic domain of CD20 and was, therefore, ascribed as CD20cy in the 5th Workshop. CD20 is a non-Ig differentiation antigen of B-cells and its expression is restricted to normal and neoplastic B-cells, being absent from all other leukocytes and tissues. CD20 is expressed by pre B-cells and persists during all stages of B-cell maturation but is lost upon terminal differentiation into plasma cells. This MAb can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin's disease, particularly of lymphocyte predominant type.

Product Info

| | |
|----------------------------|---|
| Amount : | 20 µg / 100 µg |
| Content : | 200 µg/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Storage condition : | Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. |

Application Note

Flow Cytometry (0.5-1ug/million cells); Immunofluorescence (1-2ug/ml); Western Blotting (1-2ug/ml); Immunohistology (Formalin-fixed) (0.5-1ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes),




Fig. 1: Formalin-fixed, paraffin-embedded human Tonsil stained with CD20 Mouse Monoclonal Antibody (L26).




Fig. 2: Formalin-fixed, paraffin-embedded human Lymphoma stained with CD20 Mouse Monoclonal Antibody (L26).




Fig. 3: Cytometric Analysis of Raji cells using CD20 Mouse Monoclonal Antibody (L26) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).




Fig. 4: Formalin-fixed, paraffin-embedded human Lymphoma stained with CD20 Mouse Monoclonal Antibody (L26).




Fig. 5: SDS-PAGE Analysis Purified CD20 Mouse Monoclonal Antibody (L26). Confirmation of Integrity and Purity of Antibody.




Fig. 6: Immunofluorescence staining of Raji cells using CD20 Mouse Monoclonal Antibody (L26) followed by goat anti-Mouse IgG conjugated to CF488 (green). Nuclei are stained with Reddot.




Fig. 7: Western Blot Analysis of Raji cell lysate using CD20 Mouse Monoclonal Antibody (L26).