

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

36-3560: Anti-CD28 Monoclonal Antibody(Clone: CB28)-PE

Clonality: Monoclonal

Clone Name: CB28
Application: FACS,IF
Reactivity: Human
Conjugate: PE
Gene: CD28
Gene ID: 940
Uniprot ID: P10747

Alternative Name: T-cell-specific surface glycoprotein CD28; Tp44

Isotype: Mouse IgG1, kappa

Immunogen Information: Recombinant human CD28 protein

Description

Recognizes a glycoprotein of 44-88kDa, which is identified as CD28. It is the critical T-cell co-stimulatory receptor which provides to the cell the important second activation signal by binding CD80 and CD86 that are expressed by antigen presenting cells. Besides its co-stimulation role, CD28 functions in preventing T-cells from anergic hyporesponsive state or from undergoing premature apoptotic cell death. CD28 is also expressed on human fetal NK cells and some NK cell lines, whereas on murine NK cells the CD28 expression is much broader.

Product Info

Amount: 0.5 ml at 100µg/ml

Antibody Purified from Bioreactor Concentrate by Protein A/G and conjugated to various reporter

Content: molecules. Prepared in 10mM PBS with 0.05% BSA and 0.05% azide. Contact us if you require

this Ab in a different format.

Storage condition : Antibody with azide - store at 4 to 8°C. Antibody is stable for 24 months. Non-hazardous.

Application Note

Flow Cytometry: 5ul per test per one million cells (or 5ul per 100ul of whole blood);Immunofluorescence (1:50-1:100)




Fig. 1: SDS-PAGE Analysis Purified CD28 Mouse Monoclonal Antibody (CB28). Confirmation of Integrity and Purity of Antibody.